

Horse Vaccine Comparison Chart

	Eastern Encephalomyelitis	Western Encephalomyelitis	Venezuelan Encephalomyelitis	Tetanus	Potomac Horse Fever	Rabies	Influenza	Rhinopneumonitis (equine herpesvirus)	Leptospirosis	Abortion	Endotoxemia & Diarrhea	West Nile	Strangles	Botulism Type B	Dosage	Route of Administration	Age of Initial Vaccination	First Revaccination Interval
BotVax B														•	2 ml	IM	2 wks	4 wks
Equi-Jec 6	•	•		•			•	•				•			1 ml	IM	4 mos	3-4 wks
Equi-Jec 7	•	•	•	•			•	•				•			1 ml	IM	4 mos	3-4 wks
Equi-Jec WNV												•			1 ml	IM	4 mos	3-4 wks
Equi-Jec WNV+EWT	•	•		•								•			1 ml	IM	4 mos	3-4 wks
Equiloid Innovator	•	•		•											1 ml	IM	10 mos	3-4 wks
Equine Potomavac + Imrab					•	•									1 ml	IM	3 mos	3-4 wks
EquiVac Innovator EHV-1/4								•							1 ml	IM	10 mos	3-4 wks
Fluvac Innovator							•								1 ml	IM	10 mos	3-4 wks
Fluvac Innovator 4	•	•		•			•								1 ml	IM	10 mos	3-4 wks
Fluvac Innovator 5	•	•		•			•	•							1 ml	IM	10 mos	3-4 wks
Fluvac Innovator 6	•	•	•	•			•	•							1 ml	IM	10 mos	3-4 wks
Fluvac Innovator EHV 4/1							•	•							1 ml	IM	10 mos	3-4 wks
Imrab 3						•									2 ml	IM, SQ	3 mos	1 year
IMRAB Large Animal						•									2 ml	IM, SQ	3 mos	1 year
Lepto EQ Innovator									•						1 ml	IM	6 mos	3-4 wks
Pinnacle I.N.													•	•	Single dose	IN	9 mos	2-3 wks
Pneumabort K+1b								•		•					2 ml	IM	9 mos	3-4 wks
Potomavac					•										1 ml	IM	3 mos	3-4 wks
Prestige 2							•	•							1 ml	IM	6 mos	3-4 wks
Prestige 3	•	•		•											1 ml	IM	6 mos	3-4 wks
Prestige 3 + WNV	•	•		•								•			1 ml	IM	6 mos	3-4 wks
Prestige 4	•	•		•			•								1 ml	IM	6 mos	3-4 wks
Prestige 5	•	•		•			•	•							1 ml	IM	6 mos	3-4 wks
Prestige 5 + WNV	•	•		•			•	•				•			1 ml	IM	6 mos	3-4 wks
Prestige Prodigy								•		•					2 ml	IM	6 mos	4-6 wks
Prestige V+VEE	•	•	•	•			•	•							1 ml	IM	6 mos	3-4 wks
Prestige WNV												•			1 ml	IM	6 mos	3-4 wks
Rabvac 3						•									2 ml	IM	3 mos	1 year
Strepvax II													•	•	1 ml	IM	3 mos	Twice in a Three Week Intervals
Tetanus Antitoxin				•											5 ml	IM, SQ		No Booster
Tetanus Toxoid Concentrate				•											1 ml	IM		30 days
Vetera 5 XP	•	•		•			•	•							1 ml	IM	4 mos	3-4 wks
Vetera EWT + WNV	•	•		•								•			1 ml	IM	4 mos	3-4 wks
Vetera Gold XP	•	•		•			•	•				•			1 ml	IM	4 mos	3-4 wks
Vetera VEWT + WNV	•	•	•	•								•			1 ml	IM	4 mos	3-4 wks
West Nile Innovator												•			1 ml	IM	10 mos	3-6 wks
West Nile Innovator + EWT	•	•		•								•			1 ml	IM	10 mos	3-4 wks
West Nile Innovator + VEWT	•	•	•	•								•			1 ml	IM	10 mos	3-4 wks